

**AMENDMENTS TO THE CLAIMS**

This listing of claims will replace all prior versions and listings of the claims in the application.

1. **(Currently Amended)** A method of inhibiting ~~activation proliferation~~ of a lymphocyte in a patient that may benefit from inhibition of lymphocyte activation, the method comprising administering to the patient ~~contacting an activated lymphocyte with~~ a soluble form of B7-H3 and allowing the soluble form of B7-H3 to inhibit the ~~activation proliferation~~ of the lymphocyte.
2. (Canceled)
3. (Withdrawn) The method as in claim 3, wherein the B7-H3 agonist comprises SEQ ID NO:15.
4. **(Currently Amended)** The method as in claim 1 ~~[[2]]~~, wherein the soluble form comprises at least one V domain of B7-H3.
5. **(Currently Amended)** The method as in claim 4, wherein the V domain comprises: (a) SEQ ID NO:7 or (b) an amino acid sequence which is at least 90% identical to SEQ ID NO:7 and wherein the soluble form exhibits a biological activity selected from the group consisting of decreasing T cell proliferation, decreasing secretion of IL-10, decreasing secretion of IFN- $\gamma$ , decreasing secretion of GMCSF, and decreasing secretion of TNF- $\alpha$  ~~which competitively inhibits binding of B7-H3 to its receptor~~.
6. **(Original)** The method as in claim 4, wherein the soluble form of B7-H3 further comprises at least one C domain of B7-H3.
7. **(Original)** The method as in claim 4, wherein the soluble form of B7-H3 further comprises an Fc region of an antibody.

8. **(Currently Amended)** The method as in claim 7, wherein the soluble form of B7-H3 comprises: (a) an amino acid sequence chosen from SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, or SEQ ID NO:22; or (b) an amino acid sequence which is at least 90% identical to at least one of the sequences chosen from SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, or SEQ ID NO:22 and wherein the soluble form exhibits a biological activity selected from the group consisting of decreasing T cell proliferation, decreasing secretion of IL-10, decreasing secretion of IFN- $\gamma$ , decreasing secretion of GMCSF, and decreasing secretion of TNF- $\alpha$ .~~which competitively inhibits binding of B7-H3 to its receptor.~~

9. **(Currently Amended)** The method as in claim 7, wherein the soluble form of B7-H3 comprises: (a) an amino acid sequence chosen from SEQ ID NO:10, SEQ ID NO:12, or SEQ ID NO:14; or (b) an amino acid sequence which is at least 90% identical to at least one of the sequences chosen from SEQ ID NO:10, SEQ ID NO:12, or SEQ ID NO:14 and wherein the soluble form exhibits a biological activity selected from the group consisting of decreasing T cell proliferation, decreasing secretion of IL-10, decreasing secretion of IFN- $\gamma$ , decreasing secretion of GMCSF, and decreasing secretion of TNF- $\alpha$ .~~which competitively inhibits binding of B7-H3 to its receptor.~~

10. (Withdrawn) The method as in claim 3, wherein the B7-H3 agonist is coupled with a primary stimulatory molecule.

11. (Withdrawn) The method as in claim 10, wherein the soluble form of B7-H3 and the primary stimulatory molecule are spaced by no more than 100  $\mu\text{m}$ .

12. (Withdrawn) The method as in claim 1, wherein the B7-H3 antagonist is a nucleic acid encoding amino acid of SEQ ID NO:15.

13. (Withdrawn) A method of enhancing activation of a lymphocyte, the method comprising contacting the lymphocyte with a B7-H3 antagonist and allowing the antagonist to enhance the activation of the lymphocyte.

14. (Withdrawn) The method as in claim 13, wherein the lymphocyte is human.
15. (Withdrawn) The method as in claim 13, wherein the B7-H3 antagonist is an antibody to B7-H3 or an antibody against a B7-H3 receptor.
16. (Withdrawn) The method as in claim 13, wherein the B7-H3 antagonist is an antisense nucleic acid or a siRNA.
17. **(Currently Amended)** The method as ~~in any one of~~ claim[[s]] 1, wherein the lymphocyte is a T cell.
18. **(Currently Amended)** The method as in claim[[s]] 17, wherein the T cell is a CD4<sup>+</sup> T cell.
19. (Canceled)
20. **(Currently Amended)** The method as in claim 1[[9]], wherein the patient ~~mammal~~ is afflicted with ~~or is at risk for~~ an immunologic disorder selected from the group consisting of rheumatoid arthritis, psoriasis, multiple sclerosis, inflammatory bowel disease, Crohn's disease, systemic lupus erythematosus, type I diabetes, transplant rejection, and graft-versus-host disease.
21. **(Currently Amended)** The method as in claim 19, wherein the patient ~~mammal~~ is treated with Factor VIII or Factor IX.
22. **(New)** The method as in claim 1, wherein the patient is at risk for transplant rejection or graft-versus-host disease.